

Durham Research Online

Deposited in DRO:

17 May 2017

Version of attached file:

Published Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

Israel, H. and Massey, R. and Prod'homme, T. and Cropper, M. and Cordes, O. and Gow, J. and Kohley, R. and Marggraf, O. and Niemi, S. and Rhodes, J. and Short, A. and Verhoeve, P. (2017) 'Erratum: How well can charge transfer inefficiency be corrected? A parameter sensitivity study for iterative correction.', *Monthly notices of the Royal Astronomical Society.*, 467 (4). pp. 4218-4219.

Further information on publisher's website:

<https://doi.org/10.1093/mnras/stx270>

Publisher's copyright statement:

This article has been published in *Monthly Notices of the Royal Astronomical Society* ©: 2017 The Authors Published by Oxford University Press on behalf of the Royal Astronomical Society. All rights reserved.

Additional information:

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

Erratum: How well can charge transfer inefficiency be corrected? A parameter sensitivity study for iterative correction

by Holger Israel,^{1,2,10★} Richard Massey,^{1,3★} Thibaut Prod'homme,⁴ Mark Cropper,⁵ Oliver Cordes,⁶ Jason Gow,⁷ Ralf Kohley,⁸ Ole Marggraf,⁶ Sami Niemi,⁵ Jason Rhodes,⁹ Alex Short⁴ and Peter Verhoeve⁴

¹*Institute for Computational Cosmology, Durham University, South Road, Durham DH1 3LE*

²*Centre for Extragalactic Astronomy, Durham University, South Road, Durham DH1 3LE*

³*Centre for Advanced Instrumentation, Durham University, South Road, Durham DH1 3LE*

⁴*European Space Agency, ESTEC, Keplerlaan 1, 2200AG Noordwijk, the Netherlands*

⁵*Mullard Space Science Laboratory, University College London, Holmbury St Mary, Dorking, Surrey RH5 6NT*

⁶*Argelander-Institut für Astronomie, Universität Bonn, Auf dem Hügel 71, 53121 Bonn, Germany*

⁷*e2v Centre for Electronic Imaging, The Open University, Walton Hall, Milton Keynes MK7 6AA*

⁸*European Space Agency, ESAC, PO Box 78, 28691 Villanueva de la Cañada, Madrid, Spain*

⁹*Jet Propulsion Laboratory, California Institute of Technology, 4800 Oak Grove Drive, Pasadena, CA 91109, USA*

¹⁰*Faculty of Physics, Ludwig-Maximilians-Universität, Scheinerstraße 1, 81679 München, Germany*

Key words: errata, addenda – instrumentation: detectors – methods: data analysis – space vehicles: instruments.

The article ‘How well can charge transfer inefficiency be corrected? A parameter sensitivity study for iterative correction’ was published in MNRAS 453, 561 (2015). We identified several incorrectly placed decimal points in Table 1 of that article that arose from presenting several curves in the related Fig. 2 scaled up by a factor of 10, as a visual aid. Table 1 details the parameters of a heuristic fit,

$$A + D_a \operatorname{atan}((\log \tau - D_p)/D_w) + G_a \exp(-(\log \tau - G_p)^2/2G_w^2), \quad (14)$$

to the effects caused by a species of charge traps in a CCD, created by the degradation of its silicon lattice due to striking cosmic rays outside the Earth’s atmosphere, on the measured photometry, astrometry and ellipticity of simulated galaxy images. In the published article, there is a typing error such that equation (14) lacks the negative sign in the argument of the exponential. The amended equation (14) above provides the correct form that we have been using for all fits.

The article lacks a clear notice that the effects of a single trap species given in Table 1 and Fig. 2 have to be multiplied by a factor of $2051/(464 \times 0.94) \sim 4.155$ in order to reproduce the sensitivity study presented in Section 5 (Figs 3–9) correctly. The scaling factor applies to the three amplitude parameters, i.e. A , G_a and D_a . Appendix A3 explains how the scaling factor arises when constructing the baseline trap model informed by laboratory analyses of an irradiated CCD. However, its relevance for reproducing our results using Table 1 should have been stated explicitly.

To alleviate this difficulty, we again present the coefficients of the fit to the single trap species data, but with corrected orders of magnitude and the scale factor included. We also flipped the signs of the amplitudes in the rows of Table 1 that show the results for the ellipticity offset Δe_1 , compared with the published version. The fit in Fig. 2 was made to $|\Delta e_1|$ and the new table reflects the correct sign.

Moreover, independently, we made a sign error when deriving equation (20) from the more general equation (17). Equation (20) quantifies the residual effect Δf^{Pr} on a galaxy observable if charge transfer inefficiency has been corrected using a biased or uncertain trap density, while equation (17) considers both incorrect trap densities and release time constants. The correctly derived version of equation (20) reads

$$\Delta f^{\text{Pr}}(\rho_i + \Delta \rho_i) = \sum_i \rho_i f^{\text{resid}}(\tau_i) - \sum_i \Delta \rho_i f(\tau_i), \quad (20)$$

where ρ_i and $\Delta \rho_i$ are the densities and errors in the densities of the trap species with time constants τ_i , while $f(\tau_i)$ and $f^{\text{resid}}(\tau_i)$ are the effects of a single trap species before and after correcting the charge transfer inefficiency (to be taken from the corrected Table 1 presented in this Erratum).

The correct equation (20) has been used to fit the sensitivity experiments, the findings of which are shown in Figs 5 and 9. None of the mistakes affects the results or conclusions of the original article; they are simply errors in the manuscript.

ACKNOWLEDGEMENTS

The authors thank Ruyman Azzollini, whose persistent questions about the article led to the discovery of the errors we correct here.

* E-mail: holger.israel@durham.ac.uk, hisrael@usm.lmu.de (HI); r.j.massey@durham.ac.uk (RM)

Table 1. Parameters of fitting functions to illustrate the effect on measurements of galaxy fluxes F and F_S , astrometry y and morphology R^2 , e_1 of charge traps of different species. In all cases, the measurements assume a density of one trap per pixel and the astrophysical measurement is fitted as a function of the charge trap's characteristic release time τ as $A + D_a \text{atan}((\log \tau - D_p)/D_w) + G_a \exp(-(\log \tau - G_p)^2/2G_w^2)$. Values after correction highlight the efficacy of CTI mitigation.

	A	D_a	D_p	D_w
Galaxy simulation: in degraded images, including readout noise				
$\Delta F/F_{\text{true}}$	-0.2230 ± 0.0041	-0.1307 ± 0.0035	0.620 ± 0.004	0.464 ± 0.026
Δy	0.4611 ± 0.0006	-0.2198 ± 0.0012	0.839 ± 0.008	0.211 ± 0.023
$\Delta R^2/R_{\text{true}}^2$	0.1756 ± 0.0011	-0.1603 ± 0.0016	1.572 ± 0.018	0.258 ± 0.037
Δe_1	-0.2216 ± 0.0007	0.1395 ± 0.0011	1.628 ± 0.022	0.295 ± 0.033
Galaxy simulation: after correction in software post-processing (perfect knowledge of charge traps)				
$\Delta F/F_{\text{true}}$	-0.0231 ± 0.0001	$(1.853 \pm 0.114) \times 10^{-3}$	1.296 ± 0.137	0.260 ± 0.134
Δy	$(3.981 \pm 0.420) \times 10^{-3}$	$(2.147 \pm 0.462) \times 10^{-3}$	0.56 ± 0.89	0.22 ± 0.46
$\Delta R^2/R_{\text{true}}^2$	-0.0963 ± 0.007	0.0184 ± 0.008	0.759 ± 0.250	0.285 ± 0.110
Δe_1	$(-5.743 \pm 4.758) \times 10^{-4}$	$(-1.6 \pm 2.8) \times 10^{-4}$	1.218 ± 1.932	0.100 ± 0.000
Star simulation: in degraded images, including readout noise				
$\Delta F/F_{\text{true}}$	-0.0934 ± 0.0010	-0.0605 ± 0.0008	1.075 ± 0.026	0.551 ± 0.010
Δy	0.1809 ± 0.0001	-0.0773 ± 0.0001	1.731 ± 0.004	0.292 ± 0.007
$\Delta R^2/R_{\text{true}}^2$	0.0394 ± 0.0004	-0.0267 ± 0.0004	2.888 ± 0.047	0.187 ± 0.045
Δe_1	-0.0513 ± 0.0003	0.0330 ± 0.0004	2.667 ± 0.024	0.175 ± 0.039
Star simulation: after correction in software post-processing (perfect knowledge of charge traps)				
$\Delta F/F_{\text{true}}$	$(-1.462 \pm 0.099) \times 10^{-4}$	$(1.121 \pm 0.107) \times 10^{-4}$	1.102 ± 0.105	0.422 ± 0.200
Δy	$(6.250 \pm 0.027) \times 10^{-3}$	$(4.028 \pm 0.028) \times 10^{-3}$	1.246 ± 0.186	0.273 ± 0.155
$\Delta R^2/R_{\text{true}}^2$	$(-6.784 \pm 1.593) \times 10^{-4}$	$(-7.578 \pm 1.501) \times 10^{-4}$	1.269 ± 0.329	0.25 ± 0.48
Δe_1	$(-5.084 \pm 9.972) \times 10^{-5}$	$(-1.573 \pm 5.817) \times 10^{-5}$	0.22 ± 5.09	0.100 ± 0.000
	G_a	G_p	G_w	
Galaxy simulation: in degraded images, including readout noise				
$\Delta F/F_{\text{true}}$	0.0879 ± 0.0081	4.953 ± 0.164	4.154 ± 0.239	
Δy	0.1272 ± 0.0077	0.694 ± 0.040	0.708 ± 0.021	
$\Delta R^2/R_{\text{true}}^2$	0.4515 ± 0.0186	0.438 ± 0.005	0.378 ± 0.016	
Δe_1	-0.4114 ± 0.0084	0.455 ± 0.005	0.413 ± 0.008	
Galaxy simulation: after correction in software post-processing				
$\Delta F/F_{\text{true}}$	$(5.405 \pm 0.505) \times 10^{-3}$	0.735 ± 0.068	0.568 ± 0.052	
Δy	$(3.4 \pm 4.9) \times 10^{-3}$	0.28 ± 0.54	0.32 ± 0.28	
$\Delta R^2/R_{\text{true}}^2$	0.0227 ± 0.0095	0.413 ± 0.161	0.353 ± 0.091	
Δe_1	$(-4.074 \pm 1.140) \times 10^{-3}$	0.573 ± 0.206	0.535 ± 0.205	
Star simulation: in degraded images, including readout noise				
$\Delta F/F_{\text{true}}$	0.0478 ± 0.019	4.966 ± 0.032	3.436 ± 0.044	
Δy	0.2118 ± 0.0007	0.672 ± 0.002	0.439 ± 0.002	
$\Delta R^2/R_{\text{true}}^2$	0.8408 ± 0.030	0.944 ± 0.002	0.502 ± 0.002	
Δe_1	-0.6861 ± 0.019	0.949 ± 0.002	0.526 ± 0.002	
Star simulation: after correction in software post-processing				
$\Delta F/F_{\text{true}}$	$(0.024 \pm 1.125) \times 10^{-3}$	1.826 ± 0.713	0.035 ± 1.000	
Δy	$(9.069 \pm 0.141) \times 10^{-3}$	0.738 ± 0.102	0.506 ± 0.072	
$\Delta R^2/R_{\text{true}}^2$	$(8.236 \pm 6.060) \times 10^{-4}$	0.508 ± 0.346	0.379 ± 0.386	
Δe_1	$(-1.108 \pm 0.253) \times 10^{-3}$	0.846 ± 0.180	0.538 ± 0.165	

This paper has been typeset from a \LaTeX file prepared by the author.